Faster-stronger!

A better core experience!





Dual-Cure Composite Core Build-Up Material

Build cores with confidence! CompCore™ AF resin composite is easy to place, shape and finish while its superior mechanical properties and fluoride release deliver long-term clinical success. And CompCore AF cuts like dentin with excellent visibility on radiographs.



Outstanding handling Ideal viscosity with excellent adaptation

Superior mechanical properties *Long term clinical success*



Optional dual-cure Chemical or fast light-curing in just 30 seconds

and the second second



Minimum shrinkage of < 2.0% For vital and endodontically-treated teeth

CompCore AF is Reliable to the Core!





Premier® Dental Products Company • 888-670-6100 • premierdentalco.com

Bonding Agent

(use only bonding agents compatible with dual-cure composites, such as IntegraBond™)



- 1. Clean and dry prepared tooth surface.
- 2. Apply 37% phosphoric acid to the cavosurface for 10 to 15 seconds.
- 3. Rinse and dry cavo surface, do not desiccate the tooth.
- 4. Apply first coat of IntegraBond[™] on prepared cavosurface. Allow to sit for 15 seconds, and then air dry. Apply 2 to 3 more coats; without waiting, dry again after last application.
- 5. Light-cure for 20 seconds.

Physical Properties			
Compressive Strength:	250 MPa		
Radiopacity:	400% Al		
Light-cure Setting Times			
Working Time:	2.5 minutes at 72° (22°C)		
⁺ Final Set Time	Shades		
Depth of cure	A3	White	Blue
6mm	30 sec	30 sec	30 sec
8mm	30 sec	40 sec	60 sec
Chemical-cure Setting Times			
Working Time:	2.5 minutes at 72° (22°C)		
Final set:	4.5 minutes at 98° (37°C)		

Ordering Information

CompCore[™] AF SyringeMix STACK: 9g Automix Syringes/Dual-Cure

3001512 SyringeMix STACK / Shade White

3001513 SyringeMix STACK / Shade A3

3001514 SyringeMix STACK / Shade Blue

Contains: two 9g automix syringes, 15 mix tips, 15 standard and 10 extra-fine intraoral tips.


Core Build-up







- 1. Bleed syringe. Assemble intraoral tip to mix-tip, attach to syringe and bleed tip.
- 2. Place intraoral tip into the base of the prepared tooth and dispense. Let material rise around the tip while withdrawing to eliminate air bubbles.
- 3. Light-cure for 30 seconds. If the core is larger than 8mm, lightcure for at least an additional 10 seconds.[†]
- 4. Check for complete hardness. The preparation can be finished immediately with a Two Striper[®] bur. Alternately, the core can be chemical-cured, final set-time is 4.5 minutes.



Refill Kit

3001432 Shade White3001433 Shade A3Contains: 50g cartridge of CompCore AF Dual-Cure, 25 mix tips,25 intraoral syringe tips

Accessories

3001418 SyringeMix Mix-Tips & Intraoral Tips (35 each)
3001468 SyringeMix Extra-Fine Intraoral Tips (50)
3001417 Cartridge Mix-Tips & Intraoral Tips (35 each)





Premier® Dental Products Company • 1710 Romano Drive, Plymouth Meeting, PA 19462, USA Phone 888-670-6100 • 610-239-6000 • Fax 610-239-6171 • premierdentalco.com • E-mail: info@premierdentalco.com

1. Independent laboratory study from Creighton University, data on file. Dent. Research (1153,2002) / *Two Striper is a registered trademark of the product manufacturer, Abrasive Technology, Inc.